

## Occludin (OCLN) (Tight Junctions Marker) Antibody

Mouse Monoclonal Antibody [Clone OCLN/2183]

Catalog No	Format	Size
100506658-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
100506658-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
100506658-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details

<b>Clone</b>	OCLN/2183
<b>Gene Name</b>	OCLN
<b>Immunogen</b>	Recombinant human Occludin fragment around aa 282-415 (Exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	60-82kDa
<b>Cellular Localization</b>	Cell junction, Cell membrane, Tight junction
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	HepG2 cells. Human Kidney and Pancreas



\*Optimal dilution for a specific application should be determined.

### Product Images for Occludin (OCLN) (Tight Junctions Marker) Antibody




Formalin-fixed, paraffin-embedded human Kidney stained with Occludin Mouse Monoclonal Antibody (OCLN/2183).

SDS-PAGE Analysis of Purified Occludin Mouse Monoclonal Antibody (OCLN/2183). Confirmation of Integrity and Purity of Antibody.



Western Blot Analysis of Human Pancreas and Kidney tissue lysate using Occludin Mouse Monoclonal Antibody (OCLN/2183).

Analysis of Protein Array containing more than 19,000 full-length human proteins using Occludin Monospecific Mouse Monoclonal Antibody (OCLN/2183). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human Kidney stained with Occludin Mouse Monoclonal Antibody (OCLN/2183).

### Specificity & Comments

Occludin is a tetraspan integral membrane protein in epithelial and endothelial tight junction (TJ) structures that can contain two extracellular loops. The protein exists in a variety of phosphorylated forms. Phosphorylation is involved in regulating both the localization and the function of Occludin. Expression of Occludin is upregulated by polyunsaturated fatty acids, increasing trans-endothelial cell resistance and reducing cellular permeability to large molecules. The level of Occludin varies greatly depending on tissue; in brain tissue, Occludin is highly expressed at cell-cell contact sites. Non-neural tissues show lower expression and discontinuous distribution. Up-regulation of epithelial Occludin may play a role in enhancing paracellular permeability and be related to the damage to the tight junction.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

BBB VCAM-1 Signaling, Transcription Factors