

Recombinant Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody

Rabbit Monoclonal Antibody [Clone CDH16/7028R]

Catalog No	Format	Size
1014-RBM11-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1014-RBM11-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1014-RBM11-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	CDH16/7028R
Gene Name	CDH16
Immunogen	Recombinant human full-length CDH16 protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	130kDa
Cellular Localization	Cell membrane
Species Reactivity	Human, Mouse, Rat
Positive Control	Normal kidney or renal cell carcinoma.

**Optimal dilution for a specific application should be determined.*


Product Images for Recombinant Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody



IHC analysis of formalin-fixed, paraffin-embedded human thyroid. Stained using CDH16/7028R at 2ug/ml in PBS for 30min RT. Inset: PBS instead of primary antibody; secondary only negative control.



IHC analysis of formalin-fixed, paraffin-embedded human renal cell carcinoma. Stained using CDH16/7028R at 2ug/ml in PBS for 30min RT. Inset: PBS instead of primary antibody; secondary only negative control.



SDS-PAGE Analysis of Purified CDH16 Rabbit Recombinant Monoclonal Antibody (CDH16/7028R). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

This MAb recognizes a protein of 130kDa, identified as Ksp-cadherin. Cadherins form a superfamily of related glycoproteins that mediate calcium-dependent cell adhesion and transmit signals from the extracellular matrix to the cytoplasm. Cadherins have been implicated in embryogenesis, tissue morphogenesis, tissue structure maintenance, cell polarization, neoplastic invasiveness and metastasis, and membrane transport. It is suggested that Ksp-cadherin is a marker for terminal differentiation of the basolateral membranes of renal tubular epithelial cells. Within the kidney, Ksp-Cadherin is found exclusively in the basolateral membrane of renal tubular epithelial cells and collecting duct cells, and not in glomeruli, renal interstitial cells, or blood vessels. Ksp-Cadherin has been suggested to distinguish Chromophobe Renal-Cell Carcinoma from Oncocytoma.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.