

ZBTB46 / BTBD4 / ZNF340 Antibody

Mouse Monoclonal Antibody [Clone PCR-P-ZBTB46-2B8]

| Catalog No | Format | Size |
|-------------------|---|--------|
| 140685-MSM1-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 140685-MSM1-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 140685-MSM1-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|---------------------|---|
| Flow Cytometry (Flow) | 1-2ug/million cells | |
| Immunofluorescence (IF) | 1-3ug/ml | |
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

Product Details


| | |
|-------------------------------|--|
| Clone | PCR-P-ZBTB46-2B8 |
| Gene Name | ZBTB46 |
| Immunogen | Recombinant full-length human ZBTB46 protein |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG2a |
| Mol. Weight of Antigen | 64kDa |
| Cellular Localization | Nucleus |
| Species Reactivity | Human |
| Positive Control | brain or rectum. HeLa or U-2 OS cells., Human tonsil |

*Optimal dilution for a specific application should be determined.


Product Images for ZBTB46 / BTBD4 / ZNF340 Antibody




Flow Cytometric Analysis of PFA-fixed HeLa cells. ZBTB46 Mouse Monoclonal Antibody (PCR-P-ZBTB46-2B8) followed by goat anti-mouse IgG-CF488 (blue); unstained cells (red).



Analysis of Protein Array containing more than 19,000 full-length human proteins using ZBTB46 Mouse Monoclonal Antibody (PCR-P-ZBTB46-2B8). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAB) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAB to its intended target. A MAB is considered to be specific to its intended target, if the MAB has an S-score of at least 2.5. For example, if a MAB binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAB to protein X is equal to 29.



SDS-PAGE Analysis. Purified ZBTB46 Mouse Monoclonal Antibody (PCRP-ZBTB46-2B8). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human tonsil stained with ZBTB46 Mouse Monoclonal Antibody (PCRP-ZBTB46-2B8).

Specificity & Comments

BTBD4 (BTB/POZ domain-containing protein 4), also known as zinc finger protein 340 (ZNF340) or zinc finger and BTB domain-containing protein 46 (ZBTB46), is a 589 amino acid protein that contains one BTB/POZ domain. The BTB/POZ domain mediates homomeric and heteromeric POZ-POZ interactions and is common to transcriptional regulators involved in chromatin modeling. In several BTB/POZ containing proteins, including BCL-6 and the promyelocytic leukemia zinc-finger (PLZF) oncoprotein, this domain interacts with the SMRT/N-CoR-mSin3A HDAC complex and is directly involved in repressing and silencing gene transcription. When this domain is deleted, as with the oncogenic PLZF-RAR chimera of promyelocytic leukemias, this transcriptional repression is attenuated. This suggests that BTBD4 may play a role in transcription regulation.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Dendritic Cell Marker, Nuclear Marker