

Recombinant Granulocyte-Colony Stimulating Factor (G-CSF) Antibody

Mouse Monoclonal Antibody [Clone rCSF3/900]

Catalog No	Format	Size
1440-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1440-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1440-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	rCSF3/900
Gene Name	CSF3
Immunogen	Recombinant full-length human CSF3 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	19kDa
Cellular Localization	Secreted
Species Reactivity	Human, Macaque Monkey
Positive Control	Tonsil or lymph node (IHC).

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant Granulocyte-Colony Stimulating Factor (G-CSF) Antibody



Formalin-fixed, paraffin-embedded human Tonsil stained with G-CSF Recombinant Mouse Monoclonal Antibody (rCSF3/900).



SDS-PAGE Analysis of Purified G-CSF Recombinant Mouse Monoclonal Antibody (rCSF3/900). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Tonsil stained with G-CSF Recombinant Mouse Monoclonal Antibody (rCSF3/900).

Specificity & Comments

This MAb recognizes granulocyte-colony stimulating factor (G-CSF) in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell types. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of myeloid cells. It is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes. G-CSF is a pleiotropic cytokine that influences differentiation, proliferation and activation of the neutrophilic granulocyte lineage. The human G-CSF cDNA encodes a 207 amino acid precursor containing a 29 amino acid signal peptide that is proteolytically cleaved to form a 178 amino acid residue mature protein. Two G-CSF s, which are identical except for a three amino acid deletion in the amino-terminus of one form of the protein have been isolated from human cells. Murine and human G-CSF s share 73% sequence identity at the amino acid level.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Immunology, AKT Signaling, Cytokine Signaling, Hematopoietic Stem Cells