

Desmoglein-3 (Squamous Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone 5H10]

Catalog No	Format	Size
1830-MSM4-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1830-MSM4-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1830-MSM4-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	5H10	
Gene Name	DSG3	
Immunogen	Recombinant human DSG3 extracellular protein fragment	
Host	Mouse	
Clonality	Monoclonal	
Isotype / Light Chain	IgG1 / Kappa	
Mol. Weight of Antigen	130kDa	
Cellular Localization	Cell junction, Cell membrane, Desmosome	
Species Reactivity	Human	
Positive Control	A431 cells.	

*Optimal dilution for a specific application should be determined.

Product Images for Desmoglein-3 (Squamous Cell Marker) Antibody



SDS-PAGE Analysis of Purified Desmoglein-3 Monoclonal Antibody (5H10). Confirmation of Integrity and Purity of Antibody.

Specificity & Comments

Recognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This MAb is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Desmosomes are cell-cell junctions between epithelial, myocardial, and certain other cell types. Currently, three desmoglein subfamily members are identified and all are members of the cadherin cell adhesion molecule superfamily.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Apoptosis, Autophagy, Developmental Biology

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product

2 Union Square, Union City, CA 94587, Tel: 1-866-2-NEOBIO (1-866-263-6246) orders@NeoBiotechnologies.com, www.NeoBiotechnologies.com

NeoBiotechnologies Your Partner in Proteome Research