

E2F6 Antibody

Mouse Monoclonal Antibody [Clone PCRP-E2F6-1F8]


Catalog No	Format	Size
1876-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1876-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1876-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	

Product Details	
Clone	PCRP-E2F6-1F8
Gene Name	E2F6
Immunogen	Recombinant full-length human E2F6 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	35kDa
Cellular Localization	Nucleus
Species Reactivity	Human
Positive Control	HeLa or Jurkat cells.

^{*}Optimal dilution for a specific application should be determined.


Product Images for E2F6 Antibody






Immunofluorescence Analysis of HeLa cells using E2F6 Mouse Monoclonal Antibody (PCRP-E2F6-1F8) followed by goat anti-mouse IgG-CF488 (green). Phalloidin (red).

SDS-PAGE Analysis of Purified E2F6 Mouse Monoclonal Antibody (PCRP-E2F6-1F8). Confirmation of Purity and Integrity of Antibody.



Analysis of Protein Array containing more than 19,000 full-length human proteinsusing E2F6 Mouse Monoclonal Antibody (PCRP-E2F6-1F8). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Flow Cytometric Analysis of PFA-fixed HeLa cells. E2F6 Mouse Monoclonal Antibody (PCRP-E2F6-1F8) followed by goat anti-mouse IgG-CF488 (blue); unstained cells (red).

Specificity & Comments

The human retinoblastoma gene product appears to play an important role in the negative regulation of cell proliferation. Functional inactivation of Rb can be mediated either through mutation or as a consequence of interaction with DNA tumor virus encoded proteins. Of all the Rb associations described to date, the identification of a complex between Rb and the transcription factor E2F most directly implicates Rb in regulation of cell proliferation. E2F was originally identified through its role in transcriptional activation of the adenovirus E2 promoter. Sequences homologous to the E2F binding site have been found upstream of a number of genes that encode proteins with putative functions in the G1and S phases of the cell cycle. E2F-1 is a member of a broader family of transcriptional regulators including E2F-2, E2F-3, E2F-4, E2F-5 and E2F-6, each of which forms heterodimers with a second protein, DP-1, forming an active E2F transcriptional regulatory complex.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Transcription Factors

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

