

IgA Secretory Component / ECM1 Antibody

Mouse Monoclonal Antibody [Clone ECM1/792]


Catalog No	Format	Size
1893-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1893-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1893-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	


Product Details	
Clone	ECM1/792
Gene Name	ECM1
Immunogen	Recombinant full-length human ECM1 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	~80kDa
Cellular Localization	Extracellular matrix, Extracellular space, Secreted
Species Reactivity	Human, Rat
Positive Control	Lung, or Breast tumor., Stomach

*Optimal dilution for a specific application should be determined.

Product Images for IgA Secretory Component / ECM1 Antibody



SDS-PAGE Analysis of Purified IgA Secretory Component Mouse Monoclonal Antibody (ECM1/792). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with IgA Secretory Component Mouse Monoclonal Antibody (ECM1/792).

Specificity & Comments

This MAb reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelium, and the antibody is a popular marker for identifying subpopulations of epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
