

Endoglin / CD105 (Angiogenesis Marker) Antibody

Mouse Monoclonal Antibody [Clone ENG/1327]


Catalog No	Format	Size
2022-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2022-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2022-MSM2-P1ABX	Purified Ab WITHOUT BSA at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Western Blot (WB)	2-4ug/ml	

Product Details		
Clone	ENG/1327	
Gene Name	ENG	
Immunogen	Recombinant human Endoglin protein fragment (aa74-251) (exact sequence is proprietary)	
Host	Mouse	
Clonality	Monoclonal	
Isotype / Light Chain	IgG1 / Kappa	
Mol. Weight of Antigen	84kDa	
Cellular Localization	Cell membrane	
Species Reactivity	Human	
Positive Control	A431 or HL-60 cell lysates (WB). Human colon tissue (IHC)., HeLa	

^{*}Optimal dilution for a specific application should be determined.

Product Images for Endoglin / CD105 (Angiogenesis Marker) Antibody





SDS-PAGE Analysis of Purified Endoglin / CD105 Mouse Monoclonal Antibody (ENG/1327). Confirmation of Integrity and Purity of Antibody

Western Blot of HeLa and A431 cell lysates with Endoglin / CD105 Mouse Monoclonal Antibody (ENG/1327).

Specificity & Comments

CD105/Endoglin is a Type I membrane glycoprotein located on cell surfaces and is part of the TGF-beta receptor complex. This protein has been found on endothelial cells, activated macrophages, fibroblasts, and smooth-muscle cells. Endoglin has a role in the development of the cardiovascular system and in vascular remodeling. Its expression is regulated during heart development. CD105 is highly expressed in endothelial cells during tumor angiogenesis and inflammation, with weak or negative expression in vascular endothelium of normal tissues. Angiogenesis is a promising prognostic marker in a variety of tumors. Endoglin is a more specific and sensitive marker for tumor angiogenesis than CD31 or CD34, as it labels only newly-formed blood vessels and may serve as a prognostic marker for Prostate Adenocarcinoma, and cancers of the lung, stomach, breast, and brain.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Angiogenesis, Endothelial Cell Marker, Hematopoietic Stem Cells, Mesenchymal Stem Cell Differentiation

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

