

Estrogen Receptor, alpha (Marker of Estrogen Dependence) Antibody

Mouse Monoclonal Antibody [Clone ESR1/3559]


Catalog No	Format	Size
2099-MSM29-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2099-MSM29-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2099-MSM29-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	


Product Details		
Clone	ESR1/3559	
Gene Name	ESR1	
Immunogen	Recombinant human ESR1 protein fragment (around aa129-312) (exact sequence is proprietary)	
Host	Mouse	
Clonality	Monoclonal	
Isotype / Light Chain	IgG2b / Kappa	
Mol. Weight of Antigen	~67kDa	
Cellular Localization	Cell membrane, Cytoplasm, Golgi apparatus, Nucleus	
Species Reactivity	Human	
Positive Control	T47D, MCF-7 cells. Breast or Endometrial Carcinoma (IHC).	

^{*}Optimal dilution for a specific application should be determined.


Product Images for Estrogen Receptor, alpha (Marker of Estrogen Dependence) Antibody









Flow Cytometric Analysis of MCF-7 cells using Estrogen Receptor alpha Mouse Monoclonal Antibody (ESR1/3559) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).




Analysis of Protein Array containing more than 19,000 full-length human proteinsusing Estrogen Receptor alpha Mouse Monoclonal Antibody (ESR1/3559) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD,Äôs) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD,Äôs) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Western Blot Analysis of T47D lysate using ESR1 Mouse MonoclonalAntibody (ESR1/3559).





Western Blot Analysis of T-47D lysate using Estrogen Receptor alpha Mouse Monoclonal Antibody (ESR1/3559).

SDS-PAGE Analysis of Purified Estrogen Receptor alpha Mouse Monoclonal Antibody (ESR1/3559). Confirmation of Integrity and Purity of Antibody.

Specificity & Comments

This monoclonal antibody is specific to estrogen receptor alpha (ER alpha) and shows minimal cross-reaction with other members of the family. ER is an important regulator of growth and differentiation in the mammary gland. Presence of ER in breast tumors indicates an increased likelihood of response to anti-estrogen (e.g. tamoxifen) therapy. It strongly stains nuclei of epithelial cells in breast carcinomas.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Store at 2 to 8°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Breast Cancer, Cardiovascular, Infectious Disease, Ovarian Cancer, Signal Transduction, Transcription Factors

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

