

## Fibroblast Activation Protein Alpha / FAP-1 Antibody

Mouse Monoclonal Antibody [Clone FAP/4853]

Catalog No	Format	Size
2191-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2191-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2191-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details


<b>Clone</b>	FAP/4853
<b>Gene Name</b>	FAP
<b>Immunogen</b>	Recombinant fragment (around aa 1-200) of human FAP protein (exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2c / Kappa
<b>Mol. Weight of Antigen</b>	88kDa
<b>Cellular Localization</b>	Cell membrane, Cell projection, Cell surface, Cytoplasm, Invadopodium membrane, Lamellipodium membrane, Membrane, Ruffle membrane, Secreted
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	colon or pancreas. Fibroblast specific., Human uterus

\*Optimal dilution for a specific application should be determined.



### Product Images for Fibroblast Activation Protein Alpha / FAP-1 Antibody



Formalin-fixed, paraffin-embedded human liver stained with Fibroblast Activation Protein Mouse Monoclonal Antibody (FAP/4853).



SDS-PAGE Analysis of Purified FAP Mouse Monoclonal Antibody (FAP/4853). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human placenta stained with Fibroblast Activation Protein Mouse Monoclonal Antibody (FAP/4853). Inset: PBS instead of primary antibody, secondary only negative control.

Analysis of Protein Array containing more than 19,000 full-length human proteins using Fibroblast Activation Protein Mouse Monoclonal Antibody (FAP/4853). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

### Specificity & Comments

FAP (fibroblast activation protein) is a cell surface glycoprotein and serine protease that is expressed primarily in fetal mesenchymal tissues and epithelial cancer fibroblasts. In cancer, FAP functions to promote cellular proliferation. In embryonic development, FAP functions to remodel developing tissues. FAP acts as an integral membrane gelatinase composed of N-glycosylated proteolytically inactive subunits. FAP expression on chondrocyte membranes is upregulated by the combination of the cytokines IL-1 and OSM and has been shown to increase in osteoarthritic patients. This expression is colocalized with MMP-1 and MMP-13 as well as CD44 (variants v3 and v7/8). Mice that lack all copies of the FAP gene have been found to be fertile and to have developmental defects or change in cancer susceptibility.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.