

GLG1 (Golgi Glycoprotein 1) (Marker for Human Cells) Antibody

Mouse Monoclonal Antibody [Clone GLG1/4830]

Catalog No	Format	Size
2734-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2734-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2734-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	GLG1/4830
Gene Name	GLG1
Immunogen	Recombinant full-length human GLG1 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	134kDa
Cellular Localization	Cytoplasm, Cytoskeleton, Golgi apparatus membrane, Golgi outpost, Microtubule organizing center
Species Reactivity	Human
Positive Control	A431, HePG2 or HeLa cells. Human placenta, tonsil or ovary.

*Optimal dilution for a specific application should be determined.

Product Images for GLG1 (Golgi Glycoprotein 1) (Marker for Human Cells) Antibody



SDS-PAGE Analysis of Purified GLG1 Mouse Monoclonal Antibody (GLG1/4830). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human colon stained with GLG1 Mouse Monoclonal Antibody (GLG1/4830).

Specificity & Comments

This MAb recognizes a protein of 134kDa, which binds fibroblast growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. It contains sialic acid residues and 16 Cys-rich GLG1 repeats. This MAb can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This MAb is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing through them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae. The Golgi tends to be more pronounced and numerous in cells that make and secrete many substances such as plasma B cells.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
