

GnRH-Receptor / LH-RH Receptor Antibody


Mouse Monoclonal Antibody [Clone A9E4]

Catalog No	Format	Size
2798-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2798-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2798-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note	
Product Details			
Clone	A9E4		
Gene Name	GNRHR		
Immunogen	A synthetic peptide aa 1-29 (MANSASPEQNQHCSAINNSIPLMQGNLPY) from the N-terminal of human GnRH receptor.		
Host	Mouse		
Clonality	Monoclonal		
Isotype / Light Chain	IgG1 / Kappa		
Mol. Weight of Antigen	54-60kDa		
Cellular Localization	Cell membrane		
Species Reactivity	Human, Rat		
Positive Control	ovarian or breast cancers., T47D cells. Pituitary gland		

^{*}Optimal dilution for a specific application should be determined.

Product Images for GnRH-Receptor / LH-RH Receptor Antibody



SDS-PAGE Analysis of Purified LHRH Receptor Mouse Monoclonal Antibody (A9E4). Confirmation of Purity and Integrity of Antibody.



Specificity & Comments

Recognizes an epitope on the extracellular domain of gonadotropin releasing hormone (GnRH) receptor or luteinizing hormone receptor (LHCGR). Lutropin (also designated luteinizing hormone) plays a role in spermatogenesis and ovulation by stimulating the testes and ovaries to produce steroids. Gonadotropin (also designated choriogonadotropin) production in the placenta maintains estrogen and progesterone levels during the first trimester of pregnancy. Ovaries and testes abundantly express luteinizing hormone/choriogonadotropin receptor. GnRH receptor contains seven hydrophobic transmembrane domains connected hydrophilic extracellular and intracellular loops characteristic of Gprotein coupled receptors. GnRH stimulates the gonadotrophs of the anterior pituitary to secrete luteinizing hormone (LH) as well as follicle-stimulating hormone (FSH). GnRH influences the protective effect of pregnancy and Gonadotropin against breast cancer. The expression of GnRH on breast carcinoma correlates in part to the degree of tumor differentiation. GnRH-positive breast tumors occur more frequently in tumors with greater cell differentiation in premenopausal women. GnRH is present in luteal and granulosa cells as well as in ovarian cell membrane preparations.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Signal Transduction

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

