

## Granzyme B (NK/T-Cell Lymphoma Marker) Antibody

Mouse Monoclonal Antibody [Clone GZMB/2403]

Catalog No	Format	Size
3002-MSM3-P0	Purified Ab with BSA and Azide	200ug/ml
3002-MSM3-P1	Purified Ab with BSA and Azide	200ug/ml
3002-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml



Applications	Tested Dillution

### Product Details

<b>Clone</b>	GZMB/2403
<b>Gene Name</b>	GZMB
<b>Immunogen</b>	Recombinant fragment of human GZMB protein (around aa 73-187) (exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2b / Lambda
<b>Mol. Weight of Antigen</b>	29-32kDa
<b>Cellular Localization</b>	Cytolytic granule, Secreted
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	A431 cells. Human tonsil, spleen or Hodgkin's Lymphoma tissue (IHC).

\*Optimal dilution for a specific application should be determined.

### Product Images for Granzyme B (NK/T-Cell Lymphoma Marker) Antibody



SDS-PAGE Analysis of Purified Granzyme B Mouse Monoclonal Antibody (GZMB/2403). Confirmation of Integrity and Purity of Antibody.

Analysis of Protein Array containing >19,000 full-length human proteins using Granzyme B Mouse Monoclonal Antibody (GZMB/2403) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

### **Specificity & Comments**

Granzyme B is a member of the granule serine protease family stored specifically in NK cells or cytotoxic T cells. Cytolytic T lymphocytes (CTL) and natural killer (NK) cells share the ability to recognize, bind, and lyse specific target cells. They are thought to protect their host by lysing cells bearing on their surface 'nonself' antigens, usually peptides or proteins resulting from infection by intracellular pathogens. Granzyme B is crucial for the rapid induction of target cell apoptosis by CTLs in the cell-mediated immune response. Granzyme B is useful as a marker in the identification of NK/T-cell lymphomas. High percentages of cytotoxic T-cells have been shown to be an unfavorable prognostic indicator in Hodgkin's Disease.

---

### **Research Areas**

Apoptosis, Autophagy, Cardiovascular, Immuno Oncology, Signal Transduction

---

### **Known Applications & Suggested Dilutions**

ELISA (For coating use Ab at 1-5ug/ml, order Ab without BSA)  
Optimal dilution for a specific application should be determined.

---

### **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

### **Supplied As**

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

---

### **Storage and Stability**

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---