

HLA-Aw32 & HLA-A25 (MHC I) Antibody

Mouse Monoclonal Antibody [Clone SPM417]

Catalog No	Format	Size
3105-MSM4XX-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3105-MSM4XX-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3105-MSM4XX-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	SPM417
Gene Name	HLA-B
Immunogen	Normal human peripheral blood lymphocytes of phenotype A1, Aw32, B7, B37, Cw-, Cw-, DR2, DRw10
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	kDa
Cellular Localization	Cell membrane, Endoplasmic reticulum membrane
Species Reactivity	Human
Positive Control	THP-1 and Jurkat cells. Tonsil or lymph node.

**Optimal dilution for a specific application should be determined.*

Product Images for HLA-Aw32 & HLA-A25 (MHC I) Antibody



Formalin-fixed, paraffin-embedded human Tonsil stained with MHC I Monoclonal Antibody (SPM417).

Specificity & Comments

This MAb reacts with cells bearing HLA-A25 or HLA-Aw32 antigens. In addition, a reaction was observed with a cell of phenotype A2, Aw31; B17, Bw49. HLA-A, with HLA-B and HLA-C, belongs to major histocompatibility complex (MHC) class I antigens and expresses constitutively on all nucleated cells. HLA system comprises closely linked genes controlling highly polymorphic proteins involved in the presentation of peptides to the T-cell receptor, inhibition of NK cell cytotoxicity, and rejection of tissue allotransplantation. Specific alleles at HLA loci are associated with diseases. This MAB is specifically applicable for typing peripheral T cells for the antigens HLA-A25 and HLA-Aw32.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cytokine Signaling, Dendritic Cell Marker, Immunology, Infectious Disease, Mesenchymal Stem Cell Differentiation

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
