

## HLA-DP (MHC II) Antibody

Mouse Monoclonal Antibody [Clone BRA-FB6]


Catalog No	Format	Size
3115-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3115-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3115-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	


Product Details	
<b>Clone</b>	BRA-FB6
<b>Gene Name</b>	HLA-DPB1
<b>Immunogen</b>	Non-T, non-B human acute lymphoblastic leukemia REH6 cell line
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2b / Kappa
<b>Mol. Weight of Antigen</b>	33-35kDa
<b>Cellular Localization</b>	Cell membrane, Endoplasmic reticulum membrane, Endosome membrane, Golgi apparatus, Lysosome membrane, trans-Golgi network membrane
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Raji cells. Tonsil or lymph node.

\*Optimal dilution for a specific application should be determined.

### Product Images for HLA-DP (MHC II) Antibody



Flow Cytometric Analysis of PFA-fixed Raji cells. HLA-DP Mouse Monoclonal Antibody (BRA-FB6) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).



SDS-PAGE Analysis of Purified HLA-DP Mouse Monoclonal Antibody (BRA-FB6). Confirmation of Integrity and Purity of Antibody.

### Specificity & Comments

Recognizes a non-polymorphic determinant of DP-MHC class II. MHC class II antigens are transmembrane glycoproteins of non-covalently linked alpha (33-35kDa) and beta (27-30kDa) chains. It reportedly reacts with B- non-T, non-B cell lines but not with T- and myeloid cell lines and leukemias. Differential expression of MHC class II antigens on fetal and adult lymphocytes, malignant B cells appears to reflect the stage of cell differentiation which may be useful in the study of lymphoproliferative disorders.

### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Cardiovascular, Immuno Oncology, Immunology, Cytokine Signaling, Dendritic Cell Marker

## Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---