

FOXA1 / HNF3A Antibody

Mouse Monoclonal Antibody [Clone FOXA1/1519]


Catalog No	Format	Size
3169-MSM9-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3169-MSM9-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3169-MSM9-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details		
Clone	FOXA1/1519	
Gene Name	FOXA1	
Immunogen	Recombinant fragment of human FOXA1 protein (aa372-472) (exact sequence is proprietary)	
Host	Mouse	
Clonality	Monoclonal	
Isotype / Light Chain	IgG2a / Kappa	
Mol. Weight of Antigen	49kDa	
Cellular Localization	Nucleus	
Species Reactivity	Human, Rat	
Positive Control	Bladder, colon or breast., HepG-2 or MCF-7 cells. Human liver, Pancreas, prostate	

^{*}Optimal dilution for a specific application should be determined.


Product Images for FOXA1 / HNF3A Antibody






Formalin-fixed, paraffin-embedded rat colon stained with FOXA1 Mouse Monoclonal Antibody (FOXA1/1519).


Western blot analysis of human HeLa cell lysate using FOXA1 Mouse Monoclonal Antibody (FOXA1/1519).




SDS-PAGE Analysis Purified FOXA1 Mouse Monoclonal Antibody (FOXA1/1519). Confirmation of Integrity and Purity of Antibody.



Immunofluorescent analysis of PFA-fixed MCF-7 cells labeled with FOXA1 Mouse Monoclonal Antibody (FOXA1/1519) followed by goat anti-mouse IgG-CF488. Counterstain: phalloidin.



Flow cytometric analysis of PFA-fixed MCF-7 cells using FOXA1 Mouse Monoclonal Antibody (FOXA1/1519) followed by goat anti-mouse IgG-CF488 (blue); isotype control (red).



Formalin-fixed, paraffin-embedded human prostate carcinoma stained with FOXA1 Mouse Monoclonal Antibody (FOXA1/1519).

Specificity & Comments

The transcription factor Forkhead-box A1 (FOXA1), also known as hepatocyte nuclear factor 3-alpha, is a member of the FOX class of transcription factors.HNF-1 (? and ?), HNF-3 (?, ? and ?), HNF-4 (? and ?), and HNF-6 compose, in part, a homeoprotein family designated the hepatocyte nuclear factor family. The various HNF-1 isoforms regulate transcription of genes in the liver as well as in other tissues such as kidney, small intestine and thymus. FOXA1 is expressed in normal breast ductal epithelium and other epithelium in different organs, such as lung, pancreas, bladder, prostate, and colon. Recently, FOXA1 has been shown to be a major determinant of estrogen-ER activity and endocrine response in breast cancer cells. FOXA1 expression correlates with estrogen receptor (ER)-positivity, especially in luminal subtype A breast cancers, which is associated with favorable prognosis. FOXA1 is useful in the subclassification of breast carcinomas.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Developmental Biology, Nuclear Marker, Signal Transduction, Stem Cell Differentiation

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

