

Androgen Receptor (Marker of Androgen Dependence) Antibody

Mouse Monoclonal Antibody [Clone DHTR/882]



Catalog No	Format	Size
367-MSM2-P0	Purified Ab with BSA and Azide	200ug/ml
367-MSM2-P1	Purified Ab with BSA and Azide	200ug/ml
367-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

Applications	Tested Dillution
Flow Cytometry (Flow)	1-2ug/million cells
Immunofluorescence (IF)	1-3ug/ml
Immunohistochemistry (IHC)	1-2ug/ml

Product Details	
Clone	DHTR/882
Gene Name	AR
Immunogen	Human DHTR recombinant protein fragment; aa151-350 (Exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	110kDa
Cellular Localization	Cytoplasm, Nucleus
Species Reactivity	Human
Positive Control	breast or prostate carcinoma., LNCap cells. Human testis

*Optimal dilution for a specific application should be determined.

Product Images for Androgen Receptor (Marker of Androgen Dependence) Antibody



Purified Androgen Receptor Mouse Monoclonal Antibody (DHTR/882). Confirmation of Integrity and Purity of Antibody.

Formalin-fixed, paraffin-embedded human prostate carcinoma stained with Androgen Receptor Mouse Monoclonal Antibody (DHTR/882).

Specificity & Comments

Recognizes a protein of 110kDa, which is identified as androgen receptor (AR). It reacts with full length, and the newly described A form of the receptor. It does not cross react with estrogen, progesterone, or glucocorticoid receptors. The expression of AR is reportedly inversely correlated with histologic grade i.e. well differentiated prostate tumors show higher expression than the poorly differentiated tumors. In prostate cancer, AR has been proposed, as a marker of hormone-responsiveness and thus it may be useful in identifying patients likely to benefit from anti-androgen therapy. Anti-androgen receptor has been useful clinically in differentiating morpheaform basal cell carcinoma (mBCC) from desmoplastic trichoepithelioma (DTE) in the skin. This MAb is superb for staining of formalin/paraffin tissues.

Research Areas

AKT Signaling, Signal Transduction, Transcription Factors

Known Applications & Suggested Dilutions

Flow Cytometry (1-2ug/million cells) | Immunofluorescence (1-2ug/ml) | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 1mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
