

# Cytokeratin 1 (Suprabasal Keratinocyte Marker) Antibody

Mouse Monoclonal Antibody [Clone KRT1/1840]


Catalog No	Format	Size
3848-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3848-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3848-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details		
Clone	KRT1/1840	
Gene Name	KRT1	
Immunogen	Recombinant full-length human KRT1 protein	
Host	Mouse	
Clonality	Monoclonal	
Isotype / Light Chain	IgG2a / Kappa	
Mol. Weight of Antigen	~67kDa	
Cellular Localization	Cell membrane	
Species Reactivity	Human, Rat	
Positive Control	Skin or Cervix.	

<sup>\*</sup>Optimal dilution for a specific application should be determined.


# Product Images for Cytokeratin 1 (Suprabasal Keratinocyte Marker) Antibody





Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with Cytokeratin 1 Mouse Monoclonal Antibody (KRT1/1840).

SDS-PAGE Analysis Purified Cytokeratin 1 Mouse Monoclonal Antibody (KRT1/1840). Confirmation of Integrity and Purity of Antibody.



Formalin-fixed, paraffin-embedded human Skin stained with Cytokeratin 1 Mouse Monoclonal Antibody (KRT1/1840).

#### **Specificity & Comments**

Keratins are a family of intermediate filament proteins that assemble into filaments through forming heterodimers of one type I keratin (keratins 9 to 23) and one type II keratin (keratins 1 to 8). Keratins demonstrate tissue and differentiation specific expression profiles. Cytokeratin 1 is a hetero-tetramer of two type I as well as two type II keratins, keratin-10 (KRT10). Keratin 1 is differentiation specific keratin that is one of the predominant keratins in suprabasal keratinocytes in stratified epithelia. Mutations in keratin 1 cause epidermolytic hyperkeratosis.

#### **Supplied As**

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### **Storage and Stability**

Antibody with azide - store at 2 to  $8^{\circ}$ C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

## **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.