

Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker) Antibody

Mouse Monoclonal Antibody [Clone EGP40/1372]

Catalog No	Format	Size
4072-MSM12-HRP1	Purified Ab conjugated to Horse Radish Peroxidase (HRP)	0.5 ml at 100ug/ml
4072-MSM12-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4072-MSM12-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4072-MSM12-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	EGP40/1372
Gene Name	EPCAM
Immunogen	Recombinant fragment from the extracellular domain of human EpCAM protein (around aa100-224) (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1
Mol. Weight of Antigen	40-43kDa
Cellular Localization	Cell junction, Cell surface, Lateral cell membrane, Tight junction
Species Reactivity	Human
Positive Control	MCF-7, HT29 cells or Ovarian carcinoma

**Optimal dilution for a specific application should be determined.*


Product Images for Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker) Antibody




Formalin-fixed, paraffin-embedded Human Ovarian Carcinoma stained with EpCAM Mouse Monoclonal Antibody (EGP40/1372) conjugated to HRP.




Formalin-fixed, paraffin-embedded human colorectal carcinoma stained with EpCAM-Monospecific Mouse Monoclonal Antibody (EGP40/1372).




Formalin-fixed, paraffin-embedded human hepatocellular carcinoma stained with EpCAM-Monospecific Mouse Monoclonal Antibody (EGP40/1372).





Flow Cytometric Analysis of PFA-fixed MCF-7 cells using EpCAM-Monospecific Mouse Monoclonal Antibody (EGP40/1372) followed by goat anti-mouse IgG-CF488 (blue); isotype control (red).



Immunofluorescence analysis of MCF-7 cells. EpCAM-Monospecific Mouse Monoclonal Antibody (EGP40/1372) labeled with CF488 (green). Nuclei are stained with RedDot (red).



Confocal immunofluorescence of human colorectal carcinoma. EpCAM-Monospecific Mouse Monoclonal Antibody (EGP40/1372) labeled with CF488 (green); Nuclei are labeled with Reddot (red).



Western Blot Analysis of MCF-7 lysate using EpCAM Mouse Monoclonal Antibody (EGP40/1372).

Analysis of Protein Array containing >19,000 full-length human proteins using EpCAM Mouse Monoclonal Antibody (EGP40/1372). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Specificity & Comments

EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Stem Cell Differentiation
