

Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker) Antibody

Mouse Monoclonal Antibody [Clone SPM134]

Catalog No	Format	Size
4072-MSM1X-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4072-MSM1X-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4072-MSM1X-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	SPM134
Gene Name	EPCAM
Immunogen	Small cell lung carcinoma cells
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1
Mol. Weight of Antigen	40-43kDa
Cellular Localization	Cell junction, Cell surface, Lateral cell membrane, Tight junction
Species Reactivity	Human
Positive Control	HT29 cells. Breast tumor.

*Optimal dilution for a specific application should be determined.



Product Images for Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker) Antibody



Immunofluorescence Analysis of MeOH-fixed MCF-7 cells labeling Ep-CAM with Ep-CAM Mouse Monoclonal Antibody (SPM134) followed by Goat anti-Mouse IgG-CF488 (green). The nuclear counterstain is NucSpot (Red).



Flow Cytometric Analysis of PFA fixed MCF-7 cells using EpCAM Mouse Monoclonal Antibody (SPM134) followed by Goat anti-mouse IgG-CF488 (Blue); Isotype Control (Red).



Formalin-fixed, paraffin-embedded human ovarian carcinoma stained with Ep-CAM Mouse Monoclonal Antibody (SPM134).

Specificity & Comments

EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Stem Cell Differentiation

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.