

## Recombinant Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker) Antibody

Mouse Monoclonal Antibody [Clone rVU-1D9]

Catalog No	Format	Size
4072-MSM20-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4072-MSM20-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4072-MSM20-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details


<b>Clone</b>	rVU-1D9
<b>Gene Name</b>	EPCAM
<b>Immunogen</b>	Recombinant full-length human EpCAM protein
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	40-43kDa
<b>Cellular Localization</b>	Cell junction, Lateral cell membrane, Tight junction
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	MCF-7 or HT29 cells. Breast tumor.

\*Optimal dilution for a specific application should be determined.


### Product Images for Recombinant Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker) Antibody




SDS-PAGE Analysis of Purified EpCAM Mouse Recombinant Monoclonal Antibody (rVU-1D9). Confirmation of Purity and Integrity of Antibody.



Flow Cytometric Analysis of PFA fixed MCF-7 cells using EpCAM Mouse Recombinant Monoclonal Antibody (rVU-1D9) followed by Goat anti-mouse IgG-CF488 (Blue); Isotype Control (Red).



Flow cytometric Analysis of live MCF-7 cells unstained (gray) or stained with CF488A-labeled Ep-CAM monoclonal antibody (rVU-1D9) (green)



Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with EpCAM Mouse Recombinant Monoclonal Antibody (rVU-1D9).

### Specificity & Comments

This antibody reacts with the first EGF repeat in the extracellular domain of Ep-CAM. It is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of recombinant MAbs Purified Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Stem Cell Differentiation