

## Recombinant MMP9 (Matrix Metalloproteinase 9) Antibody

Rabbit Monoclonal Antibody [Clone MMP9/2025R]

Catalog No	Format	Size
4318-RBM5-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4318-RBM5-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4318-RBM5-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details


<b>Clone</b>	MMP9/2025R
<b>Gene Name</b>	MMP9
<b>Immunogen</b>	Recombinant fragment (around aa 603-614) of human MMP-9 protein (exact sequence is proprietary)
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	92kDa
<b>Cellular Localization</b>	Extracellular matrix, Extracellular space, Secreted
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	U937 cells. Spleen or heart.

\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant MMP9 (Matrix Metalloproteinase 9) Antibody



SDS-PAGE Analysis of Purified MMP9 Rabbit Recombinant Monoclonal Antibody (MMP9/2025R).



Formalin-fixed, paraffin-embedded human spleen stained with MMP9 Rabbit Recombinant Monoclonal Antibody (MMP9/2025R).

### Specificity & Comments

The matrix metalloproteinases (MMP) are a family of peptidase enzymes responsible for the degradation of extracellular matrix components, including collagen, gelatin, fibronectin, laminin and proteoglycan. Transcription of MMP genes is differentially activated by phorbol ester, lipopolysaccharide (LPS) or staphylococcal enterotoxin B (SEB). MMP catalysis requires both calcium and zinc. MMP-9 (also designated 92 kDa type IV collagenase or gelatinase B) has been shown to degrade bone collagens in concert with MMP-1 (also designated interstitial collagenase, fibroblast collagenase or collagenase-1), and cysteine proteases and may play a role in bone osteoclastic resorption. MMP-1 is down-regulated by p53, and abnormality of p53 expression may contribute to joint degradation in rheumatoid arthritis by regulating MMP-1 expression.

2 Union Square, Union City, CA 94587, Tel: 1-866-2-NEOBIO (1-866-263-6246)  
orders@NeoBiotechnologies.com, www.NeoBiotechnologies.com

### Supplied As

200ug/ml of Ab Purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Autophagy, Cardiovascular, Developmental Biology, Immunology, Angiogenesis, BBB VCAM-1 Signaling, Bladder Cancer, Colon Cancer, Cytokine Signaling, Signal Transduction

## Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---