

MTAP (Tumor Suppressor Marker) Antibody

2-4ug/ml

Mouse Monoclonal Antibody [Clone MTAP/1813]

Catalog No	Format		Size	
4507-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml		20 ug	
4507-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml		100 ug	
4507-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml		100 ug	
Applications	Tested Dillution	Note		
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staini sections in 10mM Tr	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C	

Product Details

Western Blot (WB)

Clone	MTAP/1813		
Gene Name	MTAP		
Immunogen	Recombinant human MTAP protein fragment (aa97-196) (exact sequence is proprietary)		
Host	Mouse		
Clonality	Monoclonal		
Isotype / Light Chain	IgG2b / Kappa		
Mol. Weight of Antigen	31kDa		
Cellular Localization	Cytoplasm, Nucleus		
Species Reactivity	Human		
Positive Control	A431, HAP1 or MCF-7 cells. Kidney., HeLa, HePG2		
*Ontime! dilution for a apacific and	A431, HAP1 or MCF-7 cells. Kloney., HeLa, HePG2		

*Optimal dilution for a specific application should be determined.

Product Images for MTAP (Tumor Suppressor Marker) Antibody





followed by cooling at RT for 20 minutes

SDS-PAGE Analysis of Purified MTAP Mouse Monoclonal Antibody (MTAP/1813). Confirmation of Integrity and Purity of Antibody.

Western Blot Analysis of Human HeLa, A431, HePG2 and HAP1 cell lysate using MTAP Mouse Monoclonal Antibody (MTAP/1813).





Analysis of Protein Array containing more than 19,000 full-length human proteinsusing MTAP Mouse Monoclonal Antibody (MTAP/1813). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD,Åôs) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD,Åôs) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human Kidney stained with MTAP Mouse Monoclonal Antibody (MTAP/1813).

Specificity & Comments

Recognizes a protein of 31kDa, which is identified as MTAP (5'deoxy-5'-methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8° C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Immunology, Cytokine Signaling

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

