

Recombinant Smooth Muscle Myosin Heavy Chain (SM-MHC) (Leiomyosarcoma & Myoepithelial Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone MYH11/4337R]

Catalog No	Format	Size
4629-RBM7-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4629-RBM7-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4629-RBM7-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dilution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	MYH11/4337R
Gene Name	MYH11
Immunogen	Recombinant human fragment corresponding to Myosin residues within aa1-100 of Myosin (Smooth Muscle)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	227kDa
Cellular Localization	Melanosome
Species Reactivity	Human
Positive Control	Human smooth muscle, uterus or breast.

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant Smooth Muscle Myosin Heavy Chain (SM-MHC) (Leiomyosarcoma & Myoepithelial Cell Marker) Antibody



Formalin-fixed, paraffin-embedded human leiomyosarcoma stained with SM-MHC Recombinant Rabbit Monoclonal Antibody (MYH11/4337R).



Formalin-fixed, paraffin-embedded human leiomyosarcoma stained with SM-MHC Recombinant Rabbit Monoclonal Antibody (MYH11/4337R).

Specificity & Comments

Smooth Muscle Myosin, heavy chain (SMM-HC) is a cytoplasmic structural protein that is a major component of the contractile apparatus of the smooth muscle cells. Expression of smooth muscle myosin is developmentally regulated, appearing early and is specific for smooth muscle development. SM-MHC stains the intact myoepithelial cell (MEC) layers present in benign and in situ malignant breast and bronchioloalveolar lesions and is therefore very helpful in distinguishing between benign and malignant tumors. The antibody reacts with smooth muscle cells and myoepithelial cells, but not with myofibroblasts. It is very helpful in distinguishing between benign sclerosing breast lesions and infiltrating carcinomas in difficult cases since it strongly stains the myoepithelial layer in the benign lesions while it is negative in the infiltrating carcinomas.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Developmental Biology, Signal Transduction
