

Recombinant NKX2.2 (Neuroendocrine & Ewing's Sarcoma Marker) Antibody

Mouse Monoclonal Antibody [Clone rNX2/294]

Catalog No	Format	Size
4821-MSM6-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4821-MSM6-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4821-MSM6-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	rNX2/294
Gene Name	NKX2-2
Immunogen	Recombinant full-length human NKX2.2 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	40-50kDa
Cellular Localization	Nucleus
Species Reactivity	Chicken, Human, Mouse, Rat
Positive Control	Pancreas or Ewing's sarcoma.

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant NKX2.2 (Neuroendocrine & Ewing's Sarcoma Marker) Antibody



SDS-PAGE Analysis of Purified NKX2.2-Monospecific Recombinant Mouse Monoclonal Antibody (rNX2/294). Confirmation of Purity and Integrity of Antibody.



Analysis of Protein Array containing more than 19,000 full-length human proteins using NKX2.2-Monospecific Recombinant Mouse Monoclonal Antibody (rNX2/294) Z- and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to be specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human Ewing's Sarcoma stained with NKX2.2-Monospecific Recombinant Mouse Monoclonal Antibody (rNX2/294).

Specificity & Comments

Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein specific to Ewing sarcoma, and was shown to be differentially upregulated in Ewing sarcoma on the basis of array-based gene expression analysis. It acts as a valuable marker for Ewing sarcoma, with a sensitivity of 93% and a specificity of 89%, and aids in the differential diagnosis of small round cell tumors.

Supplied As

200ug/ml of recombinant MAb Purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Autophagy, Cardiovascular, Developmental Biology, Neuroscience, Neural Stem Cells, Stem Cell Differentiation

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
