

Perforin (Pore Forming Protein) (Apoptosis Marker) Antibody

Mouse Monoclonal Antibody [Clone SPM434]

Catalog No	Format	Size
5551-MSM1X-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5551-MSM1X-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5551-MSM1X-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note

Product Details

Clone	SPM434
Gene Name	PRF1
Immunogen	Recombinant human Perforin protein fragment (around aa 413-552) (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	75kDa
Cellular Localization	Cell membrane, Cytolytic granule, Endosome lumen, Secreted
Species Reactivity	Human
Positive Control	HDLM-2 cells. Human spleen tissue (IHC).

*Optimal dilution for a specific application should be determined.

Product Images for Perforin (Pore Forming Protein) (Apoptosis Marker) Antibody



Formalin-fixed, paraffin-embedded human Spleen stained with Perforin Mouse Monoclonal Antibody (SPM434).

Specificity & Comments

Perforin is a pore-forming protein that leads to osmotic lysis of the target cells and subsequently enables granzymes to enter the target cells and activate apoptosis. Perforin has structural and functional similarities to complement component 9 (C9). Like C9, this protein creates transmembrane tubules and is capable of lysing non-specifically a variety of target cells. It is one of the main cytolytic proteins of cytolytic granules, and is known to be a key effector molecule for T-cell- and natural killer-cell-mediated cytotoxicity. Defects in this gene cause familial hemophagocytic lymphohistiocytosis type 2 (HPLH2), a rare and lethal autosomal recessive disorder of early childhood. The expression of perforin is reportedly upregulated in activated CD8+ T-cells, natural killer cells and some CD4+ T-cells.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Apoptosis, Autophagy