

# Recombinant CD45RB (B-Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone rPTPRC/1132]

Catalog No	Format	Size
5788-MSM31-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5788-MSM31-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5788-MSM31-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details	
Clone	rPTPRC/1132
Gene Name	PTPRC
Immunogen	Recombinant full-length human CD45RB protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	180-220kDa
Cellular Localization	Cell membrane, Membrane raft
Species Reactivity	Human
Positive Control	Raji, Ramos, U-698 or GA-10 cells. Tonsil and Spleen.
*Optimal dilution for a specific application	n should be determined.

Optimal dilution for a specific application should be determined.

## Product Images for Recombinant CD45RB (B-Cell Marker) Antibody




kDa	NR	R	
250 —	-		
150 —			
100 —			
75	-		2ug loading
50	-	_	NR=Non-
37			reduced R=reduced
25 —	_	_	
20		1999	
15			
10 —		3	

Western Blot Analysis of human spleen cell lysate using CD45RB Recombinant Mouse Monoclonal Antibody (rPTPRC/1132).

SDS-PAGE Analysis of Purified CD45RB Recombinant Mouse Monoclonal Antibody (rPTPRC/1132). Confirmation of Integrity and Purity of Antibody.





Formalin-fixed, paraffin-embedded human Lymph Node stained with CD45RB Recombinant Mouse Monoclonal Antibody (rPTPRC/1132).

### **Specificity & Comments**

CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin.

#### **Supplied As**

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

#### Storage and Stability

Antibody with azide - store at 2 to  $8^{\circ}$ C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

#### **Research Areas**

B Cell Markers, Dendritic Cell Marker, Developmental Biology, Hematopoietic Stem Cells, Immunology, Mesenchymal Stem Cell Differentiation

### **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

