

Recombinant CD45RB (B-Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone PTPRC/1783R]

Catalog No	Format	Size
5788-RBM23-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5788-RBM23-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5788-RBM23-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	PTPRC/1783R
Gene Name	PTPRC
Immunogen	Recombinant human full-length CD45RB protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	180-220kDa
Cellular Localization	Cell membrane, Membrane raft
Species Reactivity	Human
Positive Control	or GA-10 cells. Human tonsil or spleen., Ramos, U-698

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant CD45RB (B-Cell Marker) Antibody




Immunofluorescent analysis of PFA-fixed Raji cells. CD45RB Rabbit Recombinant Monoclonal Antibody (PTPRC/1783R) followed by goat anti-rabbit IgG-CF488 (green). Counterstain is phalloidin (red).




Flow Cytometric Analysis of PFA-fixed Raji cells. CD45RB Rabbit Recombinant Monoclonal Antibody (PTPRC/1783R) followed by goat anti-rabbit IgG-CF488 (blue); Isotype Control (red).



Western Blot Analysis of human spleen tissue lysate using CD45RB Rabbit Recombinant Monoclonal Antibody (PTPRC/1783).



SDS-PAGE Analysis Purified CD45RB Rabbit Recombinant Monoclonal Antibody (PTPRC/1783R). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed paraffin-embedded human tonsil stained with CD45RB Rabbit Recombinant Monoclonal Antibody (PTPRC/1783R).

Specificity & Comments

CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin.

Supplied As

200ug/ml of Ab purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Developmental Biology, Immunology, B Cell Markers, Dendritic Cell Marker, Hematopoietic Stem Cells, Mesenchymal Stem Cell Differentiation

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.