

ROR-gamma / RORC (RAR-related Orphan Receptor C) Antibody

Mouse Monoclonal Antibody [Clone RORC/2941]

Catalog No	Format	Size
6097-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
6097-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
6097-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	RORC/2941
Gene Name	RORC
Immunogen	Recombinant full-length human RORC protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	63kDa
Cellular Localization	Nucleus
Species Reactivity	Human
Positive Control	MOLT4 cells. Human lymphocytes. Human liver or skeletal muscle tissue.

*Optimal dilution for a specific application should be determined.


Product Images for ROR-gamma / RORC (RAR-related Orphan Receptor C) Antibody




Formalin-fixed, paraffin-embedded human kidney stained with ROR-gamma / RORC Mouse Monoclonal Antibody (RORC/2941).




Immunofluorescence staining of PFA-fixed MOLT4 cells with ROR-gamma / RORC Mouse Monoclonal Antibody (RORC/2941) followed by goat anti-mouse IgG-CF488. Nuclei counterstained with RedDot.




Flow Cytometric Analysis of PFA-fixed MOLT4 cells using ROR-gamma /RORC Mouse Monoclonal Antibody (RORC/2941) followed by goat anti-mouse IgG-CF488 (blue); isotype control (red).




IHC analysis of formalin-fixed, paraffin-embedded human brain. Negative tissue control using RORC/2941 at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2 °: HRP-polymer, 30min. DAB, 5min.



SDS-PAGE Analysis Purified ROR-gamma / RORC Mouse Monoclonal Antibody (RORC/2941). Confirmation of Purity and Integrity of Antibody.



Analysis of Protein Array containing more than 19,000 full-length human proteins using ROR-gamma / RORC Mouse Monoclonal Antibody (RORC/2941). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human colon carcinoma stained with ROR-gamma / RORC Mouse Monoclonal Antibody (RORC/2941).

Specificity & Comments

This MAb recognizes a protein of 63kDa, identified as ROR-C. Its epitope maps in between aa1-50. The nuclear orphan receptors ROR α and ROR γ are members of the nuclear hormone receptor superfamily. Members of this family act by directly associating with DNA sequences known as hormone response elements (HREs) and typically bind DNA as either homo- or heterodimers. ROR α and ROR γ are unique in that they bind DNA as monomers. ROR α has multiple isoforms that share common DNA and putative ligand-binding domains, but differ in their amino terminal domains, which are generated by alternative RNA processing. ROR γ comprises a 560 amino acid protein that shares 50% amino acid identity with ROR α and is most highly expressed in skeletal muscle. Although these proteins are considered orphan receptors, due to a lack of defined ligands, experimental evidence has shown that melatonin may be the natural ligand for these nuclear receptors.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cytokine Signaling, Immunology, Nuclear Marker, Transcription Factors
