

Recombinant SOX2 (Embryonic Stem Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone SOX2/3169R]

| Catalog No | Format | | Size |
|----------------------------|---|---------------------|---|
| 6657-RBM3-P0 | Purified Ab with BSA and Azide at 200ug/ml | | 20 ug |
| 6657-RBM3-P1 | Purified Ab with BSA and Azide at 200ug/ml | | 100 ug |
| 6657-RBM3-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | | 100 ug |
| | | | |
| Applications | Tested Dillution | Note | |
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Stain | ing of formalin-fixed tissues requires heating tissue |


| Immunohistochemistry (IHC) | Ũ | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
|----------------------------|----------|---|
| Western Blot (WB) | 2-4ug/ml | |

Product Details


| Clone | SOX2/3169R | | |
|------------------------|---|--|--|
| Gene Name | SOX2 | | |
| Immunogen | Recombinant fragment (within aa176-305) of human SOX2 protein (exact sequence is proprietary) | | |
| Host | Rabbit | | |
| Clonality | Monoclonal | | |
| Isotype / Light Chain | IgG / Kappa | | |
| Mol. Weight of Antigen | 34kDa | | |
| Cellular Localization | Cytoplasm, Nucleus, Nucleus speckle | | |
| Species Reactivity | Dog, Human, Mouse | | |
| Positive Control | A549, NCCIT cells. Lung or Cervical carcinoma. | | |

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant SOX2 (Embryonic Stem Cell Marker) Antibody



SDS-PAGE Analysis of Purified SOX2 Recombinant Rabbit Monoclonal Antibody (SOX2/3169R). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Skin stained with SOX2Recombinant Rabbit Monoclonal Antibody (SOX2/3169R).



Specificity & Comments

SOX2 is required for stem cell maintenance in the central nervous system, and it also regulates gene expression in the stomach. SOX2 is necessary for regulating multiple transcription factors that affect Oct 3/4 expression. An essential function of SOX2 is to stabilize embryonic stem cells in a pluripotent state by maintaining the requisite level of Oct 3/4 expression. Reportedly, SOX2 is associated with aggressive phenotypes of breast, head and neck, gastric, colorectal, bladder, and small cell lung cancers. However, SOX2 is expressed in a high percentage of lung squamous cell carcinomas and has been shown to be an independent favorable prognostic marker.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8° C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Developmental Biology, Immunology, Cytokine Signaling, Neural Stem Cells, Signal Transduction, Stem Cell Differentiation, Transcription Factors

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

