

SOX4 (Master Regulator of Epithelial-Mesenchymal Transition) Antibody

Mouse Monoclonal Antibody [Clone SOX4/2540]

Catalog No	Format	Size
6659-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
6659-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
6659-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	SOX4/2540
Gene Name	SOX4
Immunogen	Recombinant full-length human SOX4 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	47kDa
Cellular Localization	Nucleus
Species Reactivity	Human
Positive Control	Breast or Colon Carcinoma.

*Optimal dilution for a specific application should be determined.



Product Images for SOX4 (Master Regulator of Epithelial-Mesenchymal Transition) Antibody



Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with SOX4 Mouse Monoclonal Antibody (SOX4/2540).



SDS-PAGE Analysis of Purified SOX4 Mouse Monoclonal Antibody (SOX4/2540). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with SOX4 Mouse Monoclonal Antibody (SOX4/2540).

Analysis of Protein Array containing more than 19,000 full-length human proteins using SOX4 Mouse Monoclonal Antibody (SOX4/2540). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Specificity & Comments

SOX4 is a member of the SOX (SRY-related HMG-box) family of transcription factors with a critical role in embryonic development and in cell-fate determination during organogenesis of the heart, pancreas, brain, and in B and T lymphocyte differentiation. SOX4 gene expression is upregulated in many cancer types, and increased SOX4 activity contributes to cellular transformation, cell survival, and metastasis. Gene expression profiling has uncovered SOX4 with upregulated activity during TGF β -induced epithelial-mesenchymal transition (EMT) in normal and cancerous breast epithelial cells. SOX4 is indispensable for EMT and cell survival *in vitro* and for primary tumor growth and metastasis *in vivo*. SOX4 is identified as a master regulator of EMT by governing the expression of the epigenetic modifier Ezh2.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Nuclear Marker, Signal Transduction