

StAR (Steroidogenic Acute Regulator) (Leydig Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone STAR/2154]


Catalog No	Format	Size
6770-MSM4-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
6770-MSM4-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
6770-MSM4-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes


Product Details	
Clone	STAR/2154
Gene Name	STAR
Immunogen	Recombinant fragment (around aa 39-108) of human STAR protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	30kDa
Cellular Localization	Mitochondrion
Species Reactivity	Human
Positive Control	K-562 cells. Human adrenal or testicular carcinoma.

*Optimal dilution for a specific application should be determined.


Product Images for StAR (Steroidogenic Acute Regulator) (Leydig Cell Marker) Antibody




Formalin-fixed, paraffin-embedded human testicular carcinoma stained with StAR Mouse Monoclonal Antibody (STAR/2154).




Formalin-fixed, paraffin-embedded human testicular carcinoma stained with StAR Mouse Monoclonal Antibody (STAR/2154).



SDS-PAGE Analysis Purified StARMouse Monoclonal Antibody (STAR/2154). Confirmation of Integrity and Purity of Antibody.



Analysis of Protein Array containing more than 19,000 full-length human proteins using StARMouse Monoclonal Antibody (STAR/2154). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human adrenal gland stained with StARMouse Monoclonal Antibody (STAR/2154).

Specificity & Comments

Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidogenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone. It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular