

## Recombinant Thyroglobulin (Thyroidal Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone rTGB/4744]



Catalog No	Format	Size
7038-MSM14-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
7038-MSM14-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
7038-MSM14-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details	
Clone	rTGB/4744
Gene Name	TG
Immunogen	Human thyroid follicular cells
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	660kDa (Dimeric Form)
Cellular Localization	Secreted
Species Reactivity	Human, Mouse, Rat
Positive Control	Thyroid

\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant Thyroglobulin (Thyroidal Cell Marker) Antibody



SDS-PAGE Analysis of Purified Thyroglobulin Mouse Recombinant Monoclonal Antibody (rTGB/4744). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human thyroid carcinoma stained with Thyroglobulin Mouse Recombinant Monoclonal Antibody (rTGB/4744).

### **Specificity & Comments**

rTGB/4744 reacts with a partially defined epitope of human thyroglobulin. This epitope is different from the epitope recognized by MAb 6E1. Thyroglobulin is a 660kDa dimeric pre-protein with multiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers undergo conformational maturation in the endoplasmic reticulum. The vast majority of follicular carcinomas of the thyroid will give positive immunoreactivity for anti-thyroglobulin even though sometimes only focally. Poorly differentiated carcinomas of the thyroid are frequently anti-thyroglobulin negative. Adenocarcinomas of other-than-thyroid origin do not react with this antibody. This antibody is useful in identification of thyroid carcinoma of the papillary and follicular types. Presence of thyroglobulin in metastatic lesions establishes the thyroid origin of tumor. Anti-thyroglobulin, combined with anti-calcitonin, can identify medullary carcinomas of the thyroid. Furthermore, anti-thyroglobulin, combined with anti-TTF1, can be a reliable marker to differentiate between primary thyroid and lung neoplasms.

---

### **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

### **Supplied As**

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

---

### **Storage and Stability**

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---