

## p53 Tumor Suppressor Protein Antibody

Mouse Monoclonal Antibody [Clone BP53-12 + DO-7]

Catalog No	Format	Size
7157-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
7157-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
7157-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details

<b>Clone</b>	BP53-12 + DO-7
<b>Gene Name</b>	TP53
<b>Immunogen</b>	Recombinant human wild-type p53 protein (BP53-12) & Recombinant human wild type p53 protein expressed in E. coli (DO-7)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2a, IgG2b / Kappa
<b>Mol. Weight of Antigen</b>	53kDa.
<b>Cellular Localization</b>	Centrosome, Cytoplasm, Cytoskeleton, Endoplasmic reticulum, Microtubule organizing center, Mitochondrion matrix, Nucleus, PML body
<b>Species Reactivity</b>	Cow, Human, Monkey
<b>Positive Control</b>	MDA-MB-231 cells. Breast or Colon carcinoma.

*\*Optimal dilution for a specific application should be determined.*

### Product Images for p53 Tumor Suppressor Protein Antibody



Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with p53 Monoclonal Antibody (BP53-12 + DO-7)

### **Specificity & Comments**

Recognizes a 53kDa protein, which is identified as p53 suppressor gene product. It reacts with the mutant as well as the wild form of p53 under denaturing and non-denaturing conditions. Its epitope maps within the N-terminus (aa 20-25) of p53 oncoprotein. p53 is a tumor suppressor gene expressed in a wide variety of tissue types and is involved in regulating cell growth, replication, and apoptosis. It binds to MDM2, SV40 T antigen and human papilloma virus E6 protein. Positive nuclear staining with p53 antibody has been reported to be a negative prognostic factor in breast carcinoma, lung carcinoma, colorectal, and urothelial carcinoma. Anti-p53 positivity has also been used to differentiate uterine serous carcinoma from endometrioid carcinoma as well as to detect intratubular germ cell neoplasia. Mutations involving p53 are found in a wide variety of malignant tumors, including breast, ovarian, bladder, colon, lung, and melanoma.

### **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### **Supplied As**

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### **Storage and Stability**

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### **Research Areas**

Apoptosis, Breast Cancer, Cancer, Cardiovascular, Hypoxia, Immunology, AKT Signaling, Bladder Cancer, Colon Cancer, Cytokine Signaling, Defective Intrinsic Apoptosis, Infectious Disease, Lung Cancer, MAPK Signaling, Nuclear Marker, Ovarian Cancer, Signal Transduction, Transcription Factors