

## GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody

Rat Monoclonal Antibody [Clone HSP90B1/1192]

Catalog No	Format	Size
7184-RTM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
7184-RTM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
7184-RTM2-P1ABX	Purified Ab WITHOUT BSA at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details

<b>Clone</b>	HSP90B1/1192
<b>Gene Name</b>	HSP90B1
<b>Immunogen</b>	Recombinant full-length human HSP90B1 protein
<b>Host</b>	Rat
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2a / Kappa
<b>Mol. Weight of Antigen</b>	94kDa
<b>Cellular Localization</b>	Endoplasmic reticulum lumen, Melanosome, Sarcoplasmic reticulum lumen
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	U20S cells. Breast Carcinoma.

\*Optimal dilution for a specific application should be determined.

### Product Images for GRP94 / HSP90B1 (Endoplasmic Reticulum Marker) Antibody



Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with GRP94 Monoclonal Antibody (HSP90B1/1192).

### Specificity & Comments

Recognizes a protein of 94kDa, which is identified as the glucose-regulated protein 94 (grp94) and also tumor rejection antigen (gp96). Grp94 shows a high degree of sequence homology with the heat shock protein 90 (hsp90). This MAb is highly specific to grp94 and shows minimal cross-reaction with other members of the HSP90 family. Grp's are a class of proteins unresponsive to heat shock and are induced by glucose deprivation. Grp94 has been briefly studied as a prognostic factor in breast cancer.

### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Cardiovascular, Hypoxia, Immunology, Cytokine Signaling

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.