

## PGP9.5 / Uchl1 Antibody

Mouse Monoclonal Antibody [Clone SPM574]

Catalog No	Format	Size
7345-MSM1X-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
7345-MSM1X-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
7345-MSM1X-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


Clone	SPM574
Gene Name	UCHL1
Immunogen	Native Uchl1 (PGP9.5) protein from brain
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	20-30kDa
Cellular Localization	Cytoplasm, Endoplasmic reticulum membrane
Species Reactivity	Cow, Human, Mouse, Pig, Rat
Positive Control	T98G cells. Cerebellum.

\*Optimal dilution for a specific application should be determined.


### Product Images for PGP9.5 / Uchl1 Antibody



Western Blot Analysis of human Brain lysate using Pgp9.5 / Uchl1 Monoclonal Antibody (SPM574).



Immunofluorescence Analysis of T98G cells labeling Pgp9.5 with Pgp9.5 / Uchl1 Mouse Monoclonal Antibody (SPM574) followed by Goat anti-Mouse IgG-CF488 (Green). The nuclear counterstain is Nucspot (Red)



Formalin-fixed, paraffin-embedded Rat Cerebellum stained with Pgp9.5 / Uchl1 Monoclonal Antibody (SPM574).

---

### Specificity & Comments

This MAb reacts with a protein of 20-30kDa, identified as PGP9.5, also known as ubiquitin carboxyl-terminal hydrolase-1 (UchL1). Initially, PGP9.5 expression in normal tissues was reported in neurons and neuroendocrine cells but later it was found in distal renal tubular epithelium, spermatogonia, Leydig cells, oocytes, melanocytes, prostatic secretory epithelium, ejaculatory duct cells, epididymis, mammary epithelial cells, Merkel cells, and dermal fibroblasts. Furthermore, immunostaining for PGP9.5 has been shown in a wide variety of mesenchymal neoplasms as well. A mutation in PGP9.5 gene is believed to cause a form of Parkinson's disease.

---

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

---

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---

### Research Areas

Neuroscience

---