

Vinculin (Marker of Age-related Macular Degeneration) Antibody

Mouse Monoclonal Antibody [Clone VCL/3617]

Catalog No	Format	Size
7414-MSM5-P0	Purified Ab with BSA and Azide	200ug/ml
7414-MSM5-P1	Purified Ab with BSA and Azide	200ug/ml
7414-MSM5-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

Applications	Tested Dillution
Flow Cytometry (Flow)	1-2ug/million cells
Immunofluorescence (IF)	1-3ug/ml
Immunohistochemistry (IHC)	1-2ug/ml

Product Details

Clone	VCL/3617			
Gene Name	VCL			
Immunogen	Recombinant full-length human Vinculin protein (VCL)			
Host	Mouse			
Clonality	Monoclonal			
Isotype / Light Chain	IgG1 / Kappa			
Mol. Weight of Antigen	125-135kDa			
Cellular Localization	Adherens junction, Cell junction, Cell membrane, Cytoplasm, Cytoskeleton, Focal adhesion, Sarcolemma			
Species Reactivity	Bird, Cow, Fish, Frog, Human, Mouse, Pig, Rabbit, Rat			
Positive Control	A431, K562, NIH3T3, THP-1 cells. Bladder and Testis, U-87			

*Optimal dilution for a specific application should be determined.

Product Images for Vinculin (Marker of Age-related Macular Degeneration) Antibody



k	Da		ĸ	NR	
250	_				
150	_			-	
100	_				
75	-	-			2ug loading
50	_	_	-		reduced
37	_	-			R=reduced
25	_	_	_		
20	-				
15	_				
10	_				

Formalin-fixed, paraffin-embedded human testis stained with VCL-Monospecific Mouse Monoclonal Antibody(VCL/3617).

SDS-PAGE Analysis VCL-Monospecific Mouse Monoclonal Antibody(VCL/3617). Confirmation of Purity and Integrity of Antibody.





Analysis of Protein Array containing more than 19,000 full-length human proteinsusing Mouse Vinculin Monoclonal Antibody (VCL/3617). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Immunofluorescence Analysis of PFA fixed U87 cells labeling VCL-Monospecific Mouse Monoclonal Antibody (VCL/3617) followed by Goat anti-mouse IgG-CF488 (Green).



Flow Cytometric Analysis of PFA-fixed U87 cells using Vinculin Mouse Monoclonal Antibody (VCL/3617) followed by goat anti-mouse IgG-CF488 (Blue); Isotype Control (Red).



Formalin-fixed, paraffin-embedded human Testicular Carcinoma stained with VCL-Monospecific Mouse Monoclonal Antibody (VCL/3617).

Specificity & Comments

This antibody recognizes both Vinculin (125kDa) and meta-Vinculin (135kDa). Focal adhesions are identified as areas within the plasma membrane of tissue culture cells that adhere tightly to the underlying substrate. In vivo, these regions are involved in the adhesion of cells to the extracellular matrix. Paxillin and vinculin are cytoskeletal, focal adhesion proteins that are components of a protein complex which links the Actin network to the plasma membrane. Vinculin binding sites have been identified on other cytoskeletal proteins, including Talin and ?-Actinin. In addition, vinculin, Talin and ?-Actinin each contain Actin binding sites. Expression of vinculin and Talin have been shown to be affected by the level of Actin expression. ?-Actinin has been shown to link Actin to integrins in the plasma membrane through interactions with the vinculin and Talin complex or by a direct interaction with integrin.

Research Areas

Cardiovascular, Immunology, Infectious Disease, Signal Transduction

Known Applications & Suggested Dilutions

Immunofluorescence (1-2ug/ml) | Flow Cytometry (1-2ug/million cells) | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8° C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.



Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

