

# FGF23 (Fibroblast Growth Factor 23) Antibody

**Tested Dillution** 

Mouse Monoclonal Antibody [Clone FGF23/638]

Catalog No	Format	Size
8074-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
8074-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
8074-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Note

## Applications

Product Details	
Clone	FGF23/638
Gene Name	FGF23
Immunogen	Recombinant human FGF23 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	12-32kDa
Cellular Localization	Secreted
Species Reactivity	Human
Positive Control	Human PBL cells. Brain tumors.

\*Optimal dilution for a specific application should be determined.

#### Product Images for FGF23 (Fibroblast Growth Factor 23) Antibody



SDS-PAGE Analysis of Purified FGF23 Mouse Monoclonal antibody (FGF23/638). Confirmation of Purity and Integrity of Antibody.

## **Specificity & Comments**

Fibroblast growth factor-1 (FGF-1), also designated acidic FGF, and fibroblast growth factor-2 (FGF-2), also designated basic FGF, are members of a family of growth factors that stimulate proliferation of cells of mesenchymal, epithelial and neuroectodermal origin. Additional members of the FGF family include the oncogenes FGF-3 (Int2) and FGF-4 (hst/Kaposi), FGF-5, FGF-6, FGF-7 (KGF), FGF-8 (AIGF), FGF-9 (GAF) and FGF-10 through FGF-23. Members of the FGF family share 30-55% amino acid sequence identity and similar gene structure, and are capable of transforming cultured cells when overexpressed in trans- fected cells. Cellular receptors for FGFs are members of a second multigene family, including four tyrosine kinases designated FIg (FGFR-1), Bek (FGFR-L), TKF and FGFR-3.

#### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

## Storage and Stability

Antibody with azide - store at 2 to  $8^{\circ}$ C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

## **Research Areas**

Breast Cancer, Cardiovascular, AKT Signaling, Infectious Disease, MAPK Signaling, Signal Transduction

#### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

