

## Nectin 4 / Nectin Cell Adhesion Molecule 4 / PVRL4 Antibody

Mouse Monoclonal Antibody [Clone NECT4/7271]

Catalog No	Format	Size
81607-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
81607-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
81607-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details

<b>Clone</b>	NECT4/7271
<b>Gene Name</b>	NECTIN4
<b>Immunogen</b>	Recombinant fragment (around aa1-200) of human NECTIN4 protein (exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2a / Kappa
<b>Mol. Weight of Antigen</b>	66kDa
<b>Cellular Localization</b>	Adherens junction, Cell junction, Cell membrane, Secreted
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Human tonsil, placenta or breast.

\*Optimal dilution for a specific application should be determined.

### Product Images for Nectin 4 / Nectin Cell Adhesion Molecule 4 / PVRL4 Antibody



Formalin-fixed, paraffin-embedded human placenta stained with Nectin 4 Mouse Monoclonal Antibody (NECT4/7271). Inset: PBS instead of primary antibody; secondary only negative control.

### **Specificity & Comments**

Homologous to the poliovirus receptor (PVR/CD155), the nectin immunoglobulin superfamily comprises four known isoforms (-1, -2, -3, and -4). The ectodomain of nectin family members comprises three Ig-like domains (V, C, C). Nectins localize at the adherens junctions (AJ) in epithelial and endothelial cells where they serve as adhesion molecules. Actin-based AJs play a role in mechanical adhesion, cellular morphogenesis and cellular differentiation. Nectin associates with the actin cytoskeleton through its interaction with the actin filament-binding protein afadin. Nectin 4 and afadin co-localize at cadherin-based adherens junctions in MDCKII epithelial cells. Nectin 4 and nectin 3 share a common binding region in the V domain of Nectin 1 and thus compete for Nectin 1 binding. The Nectin 3/4 binding domain maps to the C-C'-C''-D ? strands of the V domain of Nectin 1. Unlike other nectins, which are more widely expressed, Nectin 4 is mainly expressed in the placenta.

---

### **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

### **Supplied As**

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

---

### **Storage and Stability**

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---