

CD4 (T-Helper/Inducer Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone EDU-2]

Catalog No	Format	Size
920-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
920-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
920-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	

Product Details

Clone	EDU-2
Gene Name	CD4
Immunogen	Stimulated human leukocytes
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	55kDa
Cellular Localization	Cell membrane
Species Reactivity	Chimpanzee, Human, Macaca artcoids (Stumptail monkey), Orangutan
Positive Control	CCRF-CEM or HL-60. Lymph node or tonsil., Human peripheral blood mononuclear cells (PBMCs)

*Optimal dilution for a specific application should be determined.

Product Images for CD4 (T-Helper/Inducer Cell Marker) Antibody



SDS-PAGE Analysis of Purified CD4 Mouse Monoclonal Antibody (EDU-2).
Confirmation of Integrity and Purity of Antibody.

Specificity & Comments

Recognizes a protein of 55kDa, identified as CD4. It is a membrane glycoprotein of T lymphocytes that interacts with major histocompatibility complex class II antigens and is also a receptor for the human immunodeficiency virus. This protein is expressed not only in T lymphocytes, but also in B cells, macrophages, and granulocytes. It is also expressed in specific regions of the brain. The protein functions to initiate or augment the early phase of T-cell activation, and may function as an important mediator of indirect neuronal damage in infectious and immune-mediated diseases of the central nervous system. Multiple alternatively spliced transcript variants encoding different isoforms have been identified. This MAb was characterized as human CD4 antibody at II and IV International Workshop on Human Leukocyte Differentiation Antigens.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

AKT Signaling, Cardiovascular, Cytokine Signaling, Hematopoietic Stem Cells, Immunology, Infectious Disease

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
