

Aurora B (Proliferation Marker) Antibody

Mouse Monoclonal Antibody [Clone AURKB/1845]

Catalog No	Format		Size
9212-MSM5-P0	Purified Ab with BSA and A	zide at 200ug/ml	20 ug
9212-MSM5-P1	Purified Ab with BSA and Azide at 200ug/ml		100 ug
9212-MSM5-P1ABX	Purified Ab WITHOUT BSA		100 ug
Applications	Tested Dillution	Note	


Applications	resteu Dillution	Note
Immunohistochemistry (IHC)	5	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

D		D - 1 - 11 -
Prod	ILICT	Details


Clone	AURKB/1845	
Gene Name	AURKB	
Immunogen	Recombinant fragment (around aa 89-251) of human Aurora B protein (exact sequence is proprietary)	
Host	Mouse	
Clonality	Monoclonal	
Isotype / Light Chain	IgG1 / Kappa	
Mol. Weight of Antigen	39kDa	
Cellular Localization	Centromere, Chromosome, Cytoplasm, Cytoskeleton, Kinetochore, Midbody, Nucleus, Spindle	
Species Reactivity	Human	
Positive Control	HeLa cells. Tonsil, Hodgkin's Lymphoma or Colon Carcinoma., Placenta	
*Ontimal dilution for a apacific an	plication should be determined	

*Optimal dilution for a specific application should be determined.

Product Images for Aurora B (Proliferation Marker) Antibody



SDS-PAGE Analysis of Purified Aurora B Mouse Monoclonal Antibody (AURKB/1845). Confirmation of Integrity and Purity of Antibody.



Formalin-fixed, paraffin-embedded human Tonsil stained with Aurora B Mouse Monoclonal Antibody (AURKB/1845).

Specificity & Comments

Recognizes a protein of 39kDa, which is identified as Aurora B. The serine/threonine protein kinase aurora B (Aurora B) is a chromosomal passenger protein critical for accurate chromosome segregation, cytokinesis, protein localization to the centromere and kinetochore, correct microtubule-kinetochore attachment, and regulation of the mitotic checkpoint. Aurora B forms a tight complex with inner centrosome protein and survivin. Inactivation of any of these proteins causes similar defects in chromosome segregation. A significant overexpression of Aurora B has been found in a variety of human tumors including non-small cell lung carcinoma, astrocytoma, seminoma and carcinomas of the colon, prostate, endometrium and thyroid. The expression level of Aurora B is

associated with cell proliferation and prognosis in these tumors. 2 Union Square, Union City, CA 94587, Tel: 1-866-2-NEOBIO (1-866-263-6246) orders@NeoBiotechnologies.com, www.NeoBiotechnologies.com

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8° C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Signal Transduction, Transcription Factors



Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

