

CD14 (Monocyte / Macrophage Marker) Antibody

Mouse Monoclonal Antibody [Clone LPSR/2397]

Catalog No	Format		Size		
929-MSM7-P0	Purified Ab with BSA and Azide at 200ug/ml		20 ug		
929-MSM7-P1	Purified Ab with BSA and Azide at 200ug/ml		100 ug		
929-MSM7-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml		100 ug		
Applications	Tested Dillution	Note			
Immunohistochemistry (IHC)	1-2ua/ml	30 min at RT. Stair	30 min at RT. Staining of formalin-fixed tissues requires heating tissue		


Immunohistochemistry (IHC)	<u> </u>	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	LPSR/2397		
Gene Name	CD14		
Immunogen	Recombinant fragment of human CD14 protein (around aa 25-148) (exact sequence is proprietary)		
Host	Mouse		
Clonality	Monoclonal		
Isotype / Light Chain	IgG2b / Kappa		
Mol. Weight of Antigen	55kDa		
Cellular Localization	Cell membrane, Golgi apparatus, Membrane raft, Secreted		
Species Reactivity	Human		
Positive Control	A549, THP-1 cells and human monocytes. Lymph nodes and tonsils.		

*Optimal dilution for a specific application should be determined.

Product Images for CD14 (Monocyte / Macrophage Marker) Antibody




		R	NR	
250→				
150→			-	
100→				
75→	-			
50→	_	-		2ug loading NR=Non- reduced
37→				R=reduced
25→	-	-		
20→				
15→				
10→				


Formalin-fixed, paraffin-embedded human Tonsil stained with CD14 Mouse Monoclonal Antibody (LPSR/2397).

SDS-PAGE Analysis of Purified CD14 Mouse Monoclonal Antibody (LPSR/2397). Confirmation of Integrity and Purity of Antibody.






Western Blot Analysis of A549 cell lysate using CD14 Mouse Monoclonal Antibody (LPSR/2397).



Analysis of Protein Array containing more than 19,000 full-length human proteinsusing CD14 Mouse Monoclonal Antibody (LPSR/2397) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human Lymph Node stained with CD14 Mouse Monoclonal Antibody (LPSR/2397).

kDa TW^P 250 _____ 150 _____ 100 _____ 70 _____ 50 _____ 40 ____ 25 _____ 20 _____ 15 _____ 10 _____ 21 _____ 22 _____ 10 _____ 23 _____ 20 _____ 10 _____ 25 _____ 20 _____ 10 _____ 25 _____ 20 _____ 25 _____ 20 _____ 25 _____ 20 _____ 25 ____ 25 _____ 25 _____ 25 _____ 25 _____ 25 _____ 25 _____ 25 _____ 25 _____ 25 _____ 25 _____ 25 _____ 25 _____ 25 ____ 25 _____ 25 ____ 2

Western Blot Analysis of THP-1 cell lysate using CD14 Mouse Monoclonal Antibody (LPSR/2397).

Specificity & Comments

Recognizes a protein of 55kDa, identified as CD14 (also known lipopolysaccharide receptor). CD14 is expressed strongly on monocytes and macrophage and weakly on the surface of neutrophils. CD14 is anchored to cells by linkage to glycosylphosphatidylinositol (GPI) and functions as a high affinity receptor for complexes of LPS and LPS binding protein (LBP). Soluble CD14, also binding to LPS, acts at physiological concentration as an LPS agonist and has, at higher concentrations, an LPS antagonizing effect in cell activation.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8° C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Apoptosis, Cardiovascular, Immunology, B Cell Markers, Dendritic Cell Marker, Hematopoietic Stem Cells, Infectious Disease, MAPK Signaling, Mesenchymal Stem Cell Differentiation

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

