

CD74 (B-Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone LN-2 + CLIP/813]

Catalog No	Format	Size
972-MSM5-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
972-MSM5-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
972-MSM5-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	LN-2 + CLIP/813
Gene Name	CD74
Immunogen	SU-DHL-4 lymphoma cells (LN-2); Recombinant full-length human CD74 protein (CLIP/813)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	33-41kDa
Cellular Localization	Cell membrane, Endoplasmic reticulum membrane, Endosome, Golgi apparatus, Late endosome, Lysosome, trans-Golgi network
Species Reactivity	Human
Positive Control	Daudi or Raji cells. Tonsil or lymph node.

*Optimal dilution for a specific application should be determined.


Product Images for CD74 (B-Cell Marker) Antibody



Flow Cytometric Analysis of Raji cells. CD74 Mouse Monoclonal Antibody (LN-2+CLIP/813) followed by goat anti-Mouse IgG-CF488. Orange: cells are not permeabilized. Blue: cells are permeabilized. Isotype Control (Red).



Immunofluorescence Analysis of Raji cells labeling CD74 with CD74 Mouse Monoclonal Antibody (LN-2+ CLIP/813) followed by Goat anti-Mouse IgG-CF488 (Green). The nuclear counterstain is Reddot (Red).



Formalin-fixed, paraffin-embedded human Tonsil stained with CD74 Mouse Monoclonal Antibody (LN-2 + CLIP/813).

Specificity & Comments

It recognizes a protein of ~35kDa, identified as CD74. It is a type II transmembrane protein which binds to the peptide binding groove of newly synthesized MHC class II alpha/beta heterodimers and prevents their premature association with endogenous polypeptides. CD74 is expressed primarily by antigen presenting cells, such as B-lymphocytes (from before the pre-B cell stage to before the plasma cell stage), macrophages, and monocytes, and many epithelial cells. Anti-CD74 stains predominantly germinal center lymphocytes and B-cell lymphomas, but rarely T-cell lymphomas. Anti-CD74 has been shown to be useful in differentiating atypical fibroxanthoma (-) from malignant fibrous histiocytoma (+).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cancer, Cardiovascular, Immunology, B Cell Markers