

## Recombinant CD79a (B-Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone rIGA/764]

Catalog No	Format	Size
973-MSM11-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
973-MSM11-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
973-MSM11-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details

<b>Clone</b>	rIGA/764
<b>Gene Name</b>	CD79A
<b>Immunogen</b>	Recombinant full-length human CD79A protein
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	44kDa
<b>Cellular Localization</b>	Cell membrane
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Daudi or Ramos cells. Germinal center B- cells in a lymph node or tonsil., Raji


\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant CD79a (B-Cell Marker) Antibody




SDS-PAGE Analysis of Purified CD79a Mouse Recombinant Monoclonal Antibody (rIGA/764). Confirmation of Purity and Integrity of Antibody.


Flow Cytometric Analysis of Raji cells using CD79a Mouse Recombinant Monoclonal Antibody (rIGA/764) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).



Immunofluorescence Analysis of PFA-fixed Raji cells labeling CD79a with CD79a Mouse Recombinant Monoclonal Antibody (rIGA/764) followed by Goat anti-Mouse IgG-CF488 (Green). The nuclear counterstain is Reddot (Red)



Analysis of Protein Array containing more than 19,000 full-length human proteins using CD79a Mouse Recombinant Monoclonal Antibody (rIGA/764). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human Spleen stained with CD79a Mouse Recombinant Monoclonal Antibody (rIGA/764).

### Specificity & Comments

A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Cancer, Immunology, B Cell Markers, Hematopoietic Stem Cells, Infectious Disease, Mesenchymal Stem Cell Differentiation