

Recombinant Major Vault Protein (MVP) Antibody

Rabbit Monoclonal Antibody [Clone VP2897R]

Catalog No	Format	Size
9961-RBM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
9961-RBM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
9961-RBM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	VP2897R
Gene Name	MVP
Immunogen	Recombinant full-length human MVP protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	104-110kDa
Cellular Localization	Cytoplasm, Nuclear pore complex, Nucleus, Perinuclear region
Species Reactivity	Human
Positive Control	A549, MCF-7 or HeLa cells. Breast tumors.

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant Major Vault Protein (MVP) Antibody



Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with Major Vault Protein Rabbit Recombinant Monoclonal Antibody (VP2897R).



Western Blot Analysis of A549 lysate using Major Vault Protein (MVP) Rabbit Recombinant Monoclonal Antibody (VP2897R).



SDS-PAGE Analysis Major Vault Protein Rabbit Recombinant Monoclonal Antibody (VP2897R). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

Recognizes a protein of 104kDa-110kDa, characterized as major vault protein (MVP). Vaults are large ribonucleoprotein particles (RNPs) present in all eukaryotic cells. They have a complex morphology, including several small molecules of RNA, but a single protein species. The MVP accounts for 70% of their mass. Their shape is reminiscent of the nucleopore central plug. Treatment of cells with estradiol increases the amount of MVP in nuclear extract. The hormone-dependent interaction of vaults with ER is prevented in vitro by sodium molybdate. Antibodies to estrogen, progesterone and glucocorticoid receptors are able to co-immunoprecipitate the MVP. MVP is overexpressed in many neoplastic tissues and cell lines. Expression of MVP predicts a poor response to chemotherapy.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Immunology