

## Cig2 Antibody

Mouse Monoclonal Antibody [Clone 3A11/5]

Catalog No	Format	Size
MSM1-2827-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
MSM1-2827-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
MSM1-2827-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Western Blot (WB)	2-4ug/ml	

Product Details	
Clone	3A11/5
Gene Name	N/A
Immunogen	Bacterial S. pombe Cig2 Purified protein.
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	~50kDa
Cellular Localization	N/A
Species Reactivity	Schizosaccharomyces Pombe
Positive Control	Red blood cell lysate; Human colon and lung tissue.

\*Optimal dilution for a specific application should be determined.

### Product Images for Cig2 Antibody

#### Specificity & Comments

Cig2 is a cyclin that functions as the catalytic subunit of cyclin-dependent kinases (Cdks) in the fission yeast *Schizosaccharomyces pombe*. It is a B-type, S phase cyclin that is required for both the G1/S and G2/M cell cycle transitions and is expressed in a sharp spike that peaks during the G1/S period. Cig2 binds to Cdc2 (Cdk), and the resulting Cdc2/Cig2 complex controls the G1/S transition of the cell cycle. Disruption of Cig2 delays the onset of mitosis. The expression of the Cig2 gene is dependent on Mlu1-binding factor (MBF), and the protein is destroyed during anaphase by the APC/cyclosome (APC/C) and Skp1/Cullin-1/F-box (SCF), thereby ensuring the spike expression pattern of Cig2. SCF regulates Cig2 levels in a dual manner, transcriptionally and post-translationally, while APC/C only destroys the protein.

#### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

#### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

#### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.