

Negative Control for Mouse Monoclonal Antibodies Antibody

Mouse Monoclonal Antibody [Clone IGG1/453]


Catalog No	Format	Size
MSM2-453-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
MSM2-453-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
MSM2-453-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details		
Clone	IGG1/453	
Gene Name	N/A	
Immunogen	Recombinant full-length human IG1R protein	
Host	Mouse	
Clonality	Monoclonal	
Isotype / Light Chain	IgG1 / Kappa	
Mol. Weight of Antigen	Not Applicable	
Cellular Localization	N/A	
Species Reactivity	All species	
Positive Control	Not Applicable	
*Ontimal dilution for a anapific annihilation of		

*Optimal dilution for a specific application should be determined.

Product Images for Negative Control for Mouse Monoclonal Antibodies Antibody



Formalin-fixed, paraffin-embedded human heart stained with Negative Control for Mouse Monoclonal Antibody (IGG1/453). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

kDa	NR R	
250 — 150 — 100 — 75 — 50 —	1	
37 25 20 15 10	-	2ug loadin NR=Non- reduced R=reduced

SDS-PAGE Analysis of Purified Negative Control for Mouse Monoclonal Abs (IGG1/453). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

To serve as a true negative control for various procedures, this rabbit monoclonal IgG should be used under the identical conditions and at the same dilution as the rabbit monoclonal antibody being used for a positive reaction.

Supplied As

200ug/ml of Ab purified by Protein A Column. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MGDS required.



2 Union Square, Union City, CA 94587, Tel: 1-866-2-NEOBIO (1-866-263-6246) orders@NeoBiotechnologies.com, www.NeoBiotechnologies.com

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

