

Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker) Antibody

Mouse Monoclonal Antibody [Clone A68-B/A11]

Catalog No	Format	Size
MSM2-937-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
MSM2-937-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
MSM2-937-P1BX	Purified Ab WITHOUT BSA at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note

Product Details

Clone	A68-B/A11
Gene Name	N/A
Immunogen	Neuraminidase-treated human red blood cells
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgM / Kappa
Mol. Weight of Antigen	Multiple
Cellular Localization	N/A
Species Reactivity	Human, Mouse, Rat
Positive Control	KG1 cells. Human colorectal carcinoma tissues.

*Optimal dilution for a specific application should be determined.

Product Images for Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker) Antibody

Specificity & Comments

Recognizes a disaccharide epitope, Gal 1-3GalNAc, of Thomsen-Friedenreich (TF) antigen. It is specific for both anomeric forms of the disaccharide (TF and TF^α, including related structures on the glycolipid) and shows no cross-reactivity with sialylated glycoprotein. The Thomsen-Friedenreich antigen acts as an oncofetal antigen, with low expression in normal adult tissues but increasing to fetal levels of expression in hyperplasia or malignancy. It is considered as a pan-carcinoma marker. This MAb is capable of agglutinate desialylated red blood cells. During metastasis, the ability of malignant cells to form multicellular aggregates via homotypic or heterotypic aggregation and their adhesion to the endothelium are critical. The tumor-associated carbohydrate Thomsen-Friedenreich antigen (Gal-GalNAc) is involved in tumor cell adhesion and tissue invasion. It also causes an immune response, and overexpression of the antigen causes cancer cells to be more sensitive to natural killer cell lysis. The Thomsen-Friedenreich antigen is suppressed in normal healthy cells and represents one of the few chemically well-defined antigens associated with tumor malignancy. The presence of the Thomsen-Friedenreich antigen on the surface of cancer cells may result from a divergence from the normal pathway for O-linked glycosylation in these cells, most likely caused by inappropriate localization of the enzymes involved in synthesis of the disaccharide.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.