

HepPar-1 (Hepatocellular Marker) Antibody

Mouse Monoclonal Antibody [Clone HepPar1]

Catalog No	Format	Size
MSM4-966-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
MSM4-966-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
MSM4-966-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	HepPar1
Immunogen	Extract of a formalin-fixed, rejected-allograft of a human liver
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	Not Known
Cellular Localization	N/A
Species Reactivity	Dog, Human
Positive Control	Liver or Hepatocellular Carcinoma (HCC).

**Optimal dilution for a specific application should be determined.*

Product Images for HepPar-1 (Hepatocellular Marker) Antibody



Formalin-fixed, paraffin-embedded human Liver stained with HepPar-1 Mouse Monoclonal Antibody (HepPar1).



Formalin-fixed, paraffin-embedded human Hepatocellular Carcinoma stained with HepPar-1 Mouse Monoclonal Antibody (HepPar1).

Specificity & Comments

Hepatocyte Paraffin 1 or HepPar1 localizes to the mitochondria of hepatocytes. It is a sensitive marker for distinguishing hepatocellular carcinomas (HCC) from other metastatic carcinomas as well as cholangio-carcinomas. HCC's occur primarily in the stomach, but they are also found in many other organs. The Hepatocyte Specific Antigen may also be a useful marker for intestinal metaplasia. Reportedly, strong expression of the Hepatocyte Specific Antigen correlates with smaller tumor size and longer patient survival. Occasionally, Hepatocyte Specific Antigen is also found in gastric carcinomas as well as in a few other non-hepatic tumors.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
