

Negative Control for Rabbit Monoclonal Antibodies Antibody

Rabbit Monoclonal Antibody [Clone NCRBM/1520R]

| Catalog No | Format | Size |
|-----------------|---|--------|
| RBM1-1520-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| RBM1-1520-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| RBM1-1520-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|---------------------|---|
| Flow Cytometry (Flow) | 1-2ug/million cells | |
| Immunofluorescence (IF) | 1-3ug/ml | |
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

Product Details

| | |
|-------------------------------|--|
| Clone | NCRBM/1520R |
| Gene Name | N/A |
| Immunogen | Recombinant full-length human KRT7 protein |
| Host | Rabbit |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG / Kappa |
| Mol. Weight of Antigen | Not Applicable |
| Cellular Localization | N/A |
| Species Reactivity | All species |
| Positive Control | Not Applicable |

*Optimal dilution for a specific application should be determined.

Product Images for Negative Control for Rabbit Monoclonal Antibodies Antibody



SDS-PAGE Analysis of Purified Negative Control for Rabbit Monoclonal Abs (NCRBM/1520R). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

To serve as a true negative control for various procedures, this rabbit monoclonal IgG should be used under the identical conditions and at the same dilution as the rabbit monoclonal antibody being used for a positive reaction.

Supplied As

200ug/ml of Ab purified by Protein A Column. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

2 Union Square, Union City, CA 94587, Tel: 1-866-2-NEOBIO (1-866-263-6246)
orders@NeoBiotechnologies.com, www.NeoBiotechnologies.com